

Vaccination chart for organ transplant patient before organ transplantation

Before organ transplant, vaccine protection should be checked for vaccines listed in the table below together with a professional. It is recommended to take the vaccines before the oragan transplant and, depending on the vaccine, also after the organ transplant. More information on the recommended vaccines for use in local healthcare can be found in Finnish at the HUS website: hus.fi \rightarrow Ammattilaiselle \rightarrow Ammattilaisten palvelut ja ohjeet \rightarrow Elinsiirto ja elinluovutus \rightarrow Elinsiirtopotilaiden rokotussuositus

If the box "To be clarified" is marked, the patient should find out if they have received the vaccine and when, for example by asking their health care center or a relative for vaccine information. If the patient has received the vaccine in question or has had the disease, they mark the "Vaccine in order/had the disease" box + date and strike through the "To be clarified" box. If the patient has not received the vaccine (or has not had the disease), they mark the "Vaccine not received" box and strike through the "To be clarified" box.

After the clarification of the vaccine protection for all vaccines, those vaccines marked with "Vaccine not received" should be acquired. Physician makes the necessary prescriptions and those vaccines which the patient pays for himself/herself should be bought from the pharmacy. Those vaccines that are included in the national vaccination programme the patient get for free at the health care center. For the administration of the vaccines the patient books an appointment from for example, their own health care center or occupational health center.

Vaccine	Vaccine in order/had the disease	Vaccine not received	To be clarified	Not needed
Pneumococcal PCV20 (Apexxnar®, Prevenar 20®)				
Meningococcal ACW135Y (Menveo® tai Nimenrix®)				
Meningococcal B (Bexero®, Trumemba®)				
Haemophilus influenzae-B (HIB) (ACT-HIB®, Hiberix®)				
Diphtheria-tetanus (dT) (Ditebooster®) or Diphtheria-tetanus and pertussis (dtap) (Triaxis®, Boostrix®, Ditekibooster®)				
Influenza (Vaxigriptetra®)				
Hepatit A ja B (Twinrix®) Hepatit A (Havrix®) Hepatit B (Engerix-b®, HBVAXPRO®)				
Chicken pox (Varilrix®)				
Human papillomavirus, HPV (Gardasil®)				
Measles, mumps and rubella vaccine (MMR) (Priorix®, M-M-RVAXPRO®)				
For COVID-19 vaccines, the current recommendations for risk groups from THL (Finnish Institute for Health and Welfare) are followed	E			